



# Opioid Addiction Treatment

A Guide for Patients, Families and Friends



**ASAM** American Society of  
Addiction Medicine

# INTRODUCTION

Addiction is a chronic medical illness that affects millions of people all over the world, as well as their families and communities. In 2018, about 2.1 million Americans ages 12 or older had an opioid use disorder (an active opioid addiction). Like diabetes or heart disease, there is no cure for addiction. But it can be **managed**, and people with addiction can and do recover.

There are treatments available that are proven to be safe and effective. Treatment provided by trained clinicians can lead to a healthier way of life, free from the symptoms of addiction. This healthy way of life is referred to as **recovery**. Treatment with a medication along with counseling and other support is often the most effective choice for opioid addiction.

This document provides facts about treatment from The American Society of Addiction Medicine (ASAM) — the leading medical society for addiction treatment. Learn more about ASAM at [www.ASAM.org](http://www.ASAM.org).

## DEFINITION OF ADDICTION\*

**Addiction is a treatable, chronic medical disease in which a person compulsively seeks and uses substances like drugs or alcohol, or engages in other behaviors (such as gambling) despite the harms that it causes to their health and/or their life. It is a brain disease because addiction changes how the brain works. Addiction impacts areas of the brain that control motivation, impulse control, reaction to stress, memory, and decision-making. These changes may last a long time, even after the person stops using substances. Without treatment and recovery support, addiction may keep getting worse.**

\*Modified from ASAM Definition of Addiction  
<https://www.asam.org/quality-care/definition-of-addiction>

## ASSESSMENT

Seeking help is the first important step to recovery. The next step in the process is to meet with a qualified clinician. A clinician is a health professional such as a physician, psychiatrist, psychologist, social worker, or nurse. During this first appointment, the clinician will screen the patient for an opioid use disorder and complete an assessment. The goal of the assessment is to gain a thorough understanding of patients and the severity of their illness. This will help the clinician and patient develop a treatment plan that best matches the patient's needs.

### Getting Started

- The clinician will ask questions to understand all of the factors that contribute to the patient's addiction. The more that is known, the better treatment can be planned with the patient. The patient will likely be asked about:
  - Current drug and alcohol use
  - Withdrawal symptoms
  - History of substance use
  - Family history of addiction
  - Mental or physical health problems
  - Support system
  - Available safe, stable housing
  - Motivation to change
- The next step is a complete physical examination to check the patient's overall health. This includes finding other common conditions (physical or mental) related to addiction that may impact the plan for treatment.
- The physical examination will often include a drug test.



## TREATMENT OVERVIEW

During the assessment, the clinician will also identify the patient's psychological and social support needs. They will likely recommend counseling or recovery support services, such as mutual support group participation. However, participation in counseling and other support services should not be required to receive treatment with medication. If the patient does participate in mutual support groups it is important to find a group that is supportive of the use of medications for opioid use disorder.

While a comprehensive assessment and medical evaluation are important for developing a treatment plan, completion of all assessments should not delay or prevent the patient from starting medication.

After the assessment, the clinician will discuss all recommended treatment options with the patient. Every patient situation is different, so choosing the best options is a shared decision between the patient and the clinician.

There are three main choices for medication to treat opioid addiction: methadone, buprenorphine, and naltrexone. These medications are typically recommended in combination with counseling and other support services.

There are a range of different treatment options depending on the severity of the patient's illness and other factors. ASAM describes four broad levels of care:

- ASAM Level 1 - Outpatient treatment. For example, treatment with buprenorphine or naltrexone with a primary care provider.
- ASAM Level 2 - Intensive outpatient or partial hospitalization. In IOP and PHP programs the patient receives over 9 or 20 hours per week of treatment services, respectively.
- ASAM Level 3 - Residential treatment. Residential treatment programs provide treatment in a safe, homelike setting where the patient can practice skills needed for recovery.
- ASAM Level 4 - Hospital inpatient treatment. Inpatient treatment may be needed for patients with severe medical or psychiatric problems that require medical and nurse care.

It's important to discuss the many different treatment options with the clinician to determine which is most appropriate.

## Treatment Plan

- After discussing the assessment results and treatment choices with the clinician, it's time to develop the full treatment plan.
- It is common for both the patient and clinician to sign an agreement about what to expect during treatment. This can include:
  - Treatment goals
  - Medications: usually methadone, buprenorphine or naltrexone for opioid use disorder
  - Treatment schedule
  - Counseling plan
  - Recovery support services
  - Patient commitment to cooperate with treatment
  - Risks of relapse and other safety concerns

## Patient Participation

- Patient engagement in treatment and recovery improves outcomes. Treatment should be long term, since addiction is a chronic disease. For this reason, it is important that patients work with the clinicians to develop a treatment plan that is tailored to their needs and goals.
- Patients should expect to be treated with respect and dignity and have their concerns listened to when starting or changing the treatment plan.
- To avoid health problems, patients must let their clinicians know about any other medications they are taking or if they drink alcohol. This is very important — certain medications and alcohol can cause major problems with treatment medications.
- Common patient responsibilities include:
  - Keeping all appointments
  - Providing samples for drug testing on a regular basis
  - Taking medications as prescribed
  - Using only medications that are prescribed
  - Allowing and encouraging involvement of supportive family and friends
  - Avoiding persons, places and situations that may put them at risk of **relapse**.

## Counseling

- Counseling is an important part of treatment for many patients and is usually recommended with medications.
- Counseling should be provided by a qualified clinician. Patients may receive counseling in the same place where they receive medication or in another location.
- Counseling helps patients address personal, social or other problems that may contribute to their addiction. Examples can be:
  - Increasing motivation for treatment and recovery
  - Difficult situations at work or home
  - Building a support system of people who are supportive of recovery
- All of the clinicians involved in patient treatment should work together to coordinate care.
- In addition to counseling, patients are encouraged to join support groups that include others who are also in recovery.
- Counseling and support groups are not limited to the patient. There are also many counseling services and support groups for family and friends of those dealing with addiction. Remember, addiction affects friends and family as well.

## Support from Family and Friends

- Families and friends can play a key role in supporting their loved ones to achieve a long-lasting recovery.
- It is very important that patients avoid the people, places and other things that trigger their desire to use alcohol or drugs. They also need to learn how to respond to those triggers without using alcohol or drugs. Friends and family can provide support and encouragement throughout this process.



## WITHDRAWAL

- Opioid **withdrawal** refers to the wide range of painful and unpleasant symptoms that occur after stopping or cutting down on the use of opioids. Withdrawal can last more than 10 days but is most often between 3–5 days.
- Although it can cause very troubling symptoms (such as vomiting, cramps and sweating), opioid withdrawal is rarely life-threatening.
- Using medications to control withdrawal (also called **withdrawal management**) is almost always recommended over trying to endure withdrawal without treatment. When patients do try to stop using opioids without treatment it can lead to stronger cravings and continued use.
- Withdrawal management on its own is not a treatment for opioid addiction and may increase the risk of relapse, overdose, and overdose death.
- When treating someone for opioid addiction, medication (methadone, buprenorphine, or naltrexone) in combination with counseling and other support is usually recommended.
- Lofexidine and clonidine are safe and effective for management of opioid withdrawal, but methadone and buprenorphine are more effective and can be continued for the treatment of opioid addiction.
- Opioid withdrawal management using ultra-rapid opioid detoxification (UROD) is not recommended due to high risk for adverse events including death.

## MEDICATIONS TO TREAT OPIOID ADDICTION

| Medication                  | Brand Names   |
|-----------------------------|---|
| Buprenorphine               | Sublocade® (Injection), Brixadi® (injection), generics (films or tablets) |
| Buprenorphine and naloxone  | Bunavail®, Cassipa®, Suboxone®, Zubsolv®, generics (films or tablets)     |
| Methadone                   | generics (liquid or tablets to dissolve in water)                         |
| Extended release naltrexone | Vivitrol® (injection)   |

## OPIOID ADDICTION MEDICATION

Methadone and buprenorphine are the most effective treatments available for active opioid addiction. Naltrexone is an effective treatment for preventing a relapse in patients who are no longer physically dependent on opioids. Most treatment plans will include medication. The type of medication chosen depends on a number of factors including the severity of the patient's illness, any co-occurring illnesses, access to different forms of treatment, and individual preferences.

- All FDA-approved medications—including methadone, buprenorphine and naltrexone—for the treatment of opioid use disorder should be available to all patients.
- Counseling is recommended with the use of any of these medications. However, counseling should not be required in order to receive treatment with medication.
- Patient preferences, history, and illness severity should all be considered when deciding which medication is best for each patient.
- Each medication works in a different way and has its own risks and benefits. Once started, these medications can be safely taken for years. The clinician and the patient should review together the risks and benefits of each medication.
- When used properly, these medications help restore balance to the parts of the brain that are impacted by addiction, helping patients to manage their addiction so they can recover.
- There is no typical timeline for use of medication. It differs for each patient. Some patients may need medications for many years.
- Some people have special situations such as pregnancy, mental health issues, pain, or criminal justice system involvement. They should discuss these issues with their clinician to find the right medication for their needs.
- Adolescents and their caregivers should also discuss medication options with their clinicians.

**If the first medication selected does not work well, the patient should discuss with the clinician a change to a different medication that may be more effective for addressing the patient's needs.**



## Buprenorphine

- Buprenorphine acts on the same parts of the brain as opioid medications or drugs. It helps to stabilize them to reduce the desire to use opioids and to prevent withdrawal symptoms.
- The FDA recently approved several new buprenorphine formulations for treatment of opioid use disorder.
- Buprenorphine comes as tablets or films (for daily use), extended-release injections (weekly or monthly) and an extended-release implant that is placed under the skin (every 6 months). The dosage may be adjusted over time to find the dose that works best for the patient.
- Many versions of this medication are combined with naloxone to prevent possible misuse. *Learn more about naloxone on page 12.* When misused (injected, snorted, or other) the naloxone in the buprenorphine medication can bring on withdrawal symptoms.
- Insurance coverage and price may help determine which form should be chosen. The clinician will make sure the patient has a dosage and form that meets the patient's medical needs.
- Patients may need to see their clinician frequently at the beginning of treatment, until their addiction symptoms have stabilized.
- Buprenorphine treatment can be started either at home or in a doctor's office. Both options are considered safe and effective when starting.
- Patients should wait until they are experiencing mild to moderate opioid withdrawal symptoms before taking the first dose of buprenorphine. If buprenorphine is taken too soon, it can bring on withdrawal symptoms instead of relieving them.
- Many patients will need to continue taking buprenorphine for an extended period of time. Stopping buprenorphine can put the patient at risk for relapse and overdose. This should only be done slowly with careful monitoring by the clinician.
- Not all clinicians can prescribe buprenorphine, so it is important to find an approved clinician who is able to prescribe this medication.
- Treatment with buprenorphine has been shown to reduce the patient's risk for opioid overdose death.

## Methadone

- Methadone also acts on the same part of the brain as opioid medications or drugs and helps to stabilize them to reduce the desire to use opioids and to prevent withdrawal symptoms.
- Methadone can be safely started before the patient starts to experience withdrawal symptoms.
- The clinician will work closely with the patient to find the right dose that reduces cravings without causing the patient to feel drowsy or sedated.
- Methadone comes as a liquid, or as a powder or tablet that is dissolved in liquid before it is taken. It is typically taken once per day (this may be different for patients who are pregnant or who are being treated for pain).
- Only specialized Opioid Treatment Programs are allowed to treat patients with methadone.
- At the beginning of treatment the patients will typically need to go to the clinic every day to take their daily dose.
- People who are in stable recovery may be provided a supply of medication to take at home.
- The administration of methadone should be monitored because unsupervised administration can lead to misuse and diversion.
- Many patients will need to continue taking methadone for an extended period of time. Stopping methadone can put the patient at risk for relapse and overdose. This should only be done slowly with careful monitoring by the clinician.
- Treatment with methadone has been shown to reduce the patient's risk for opioid overdose death.

## Naltrexone

- Naltrexone works by blocking opioids from acting on the brain. This means that the patient won't get high from using opioids while taking naltrexone. Naltrexone is a good option for preventing relapse in patients who are no longer dependent on opioids.
- Patients cannot have any opioids in their body when starting naltrexone. If they do, it will quickly cause severe withdrawal. Before a clinician starts a patient with an active opioid addiction on naltrexone, the patient must go through withdrawal under the clinician's supervision. This period can last anywhere from 6–10 days.
- Naltrexone is given in an extended release form that is injected in the buttocks. The injection is administered by the clinician in the office once every 3–4 weeks.
- Naltrexone is also available as a pill, but this form is **NOT** recommended except under very limited circumstances.
- Many patients will need to continue taking naltrexone for an extended period of time. Because naltrexone does not produce physical dependence, it can be stopped abruptly without withdrawal symptoms. However, stopping naltrexone can put the patient at risk for relapse and overdose. This should only be done with careful monitoring by the clinician.

Rx

PATIENT NAME \_\_\_\_\_

ADDRESS \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_

Prescription:

## **NALOXONE**

**Naloxone is used to treat an opioid overdose (it works only for overdoses from opioids). It is available as a nasal spray and in an injectable form.**

**Having naloxone available can help to save a person who has taken so much opioid that they stop or have trouble breathing.**

**If an overdose is suspected, call 911 immediately and give naloxone as soon as possible.**

**More than one dose of naloxone may be needed to reverse an overdose of high potency opioids, such as fentanyl.**

**Naloxone may need to be given multiple times since its effects may wear off before the opioid does.**

**To ensure patient safety, patients, their family members, significant others, and friends should keep naloxone on hand and should be trained in its use in case of overdose.**

**The clinician should provide naloxone, or a prescription for naloxone, for both the patient and family members. In most states, naloxone is also available without a prescription through the pharmacy.**

**After patients overdose it is important that they either begin treatment for opioid addiction or work with their clinician to adjust their treatment plan.**

## HOW DO THESE MEDICATIONS WORK?

### Methadone

*Long-lasting effect*

- **Effect:** Acts on the same areas of the brain as opioid medications or drugs but with longer lasting effects to stabilize activity in these areas
- **Use:** Treatment of opioid withdrawal and long-term treatment for opioid addiction
- **PROs:** Works well to prevent withdrawal, cravings, and use of illicit opioids; reduces risk for opioid overdose death
- **CONs:**
  - » Frequent clinic visits
  - » Can cause overdose if not taken as directed by the clinician

### Buprenorphine

*Often combined with naloxone*

- **Effect:** Acts on the same areas of the brain as opioid medications or drugs but with longer lasting effects to stabilize activity in these areas
- **Use:** Treatment of opioid withdrawal and long-term treatment for opioid addiction
- **PROs:** Works well to prevent withdrawal, cravings, and use of illicit opioids; reduces risk for opioid overdose death
- **CONs:** Risk for overdose when combined with alcohol or sedatives

### Naltrexone

- **Effect:** Blocks effect of opioids
- **Use:** Prevent relapse in patients who are no longer dependent on opioids
- **PROs:** No risk for dependence; reduces cravings
- **CONs:** Patients are more likely to drop out of treatment with naltrexone

### Naloxone

*Short-acting effect*

- **Effect:** Blocks effect of opioids
- **Use:** Treats opioid overdose
- **PROs:** Saves lives
- **CONs:** N/A

## Medication Use Summary

- Every patient is different. The right medication allows the patient to feel normal, have few side effects, feel no withdrawal symptoms, and have cravings under control.
- When used correctly, and when the treatment plan is being followed, these medications can be taken safely for years.
- Any plans to stop taking a medication, change dosage, or switch medications should always be discussed with the clinician. This should be a shared decision, with all benefits and risks discussed and understood by the patient.
- Patients will often participate in counseling and support groups based on a treatment plan agreed upon by the patient and the clinician. However, participation in counseling and other support services should not be required to receive treatment with medication.
- Relapse may occur as part of this chronic disease. If a patient relapses while using medication, the clinician will revise the treatment plan and treatment goals as needed.
- Naloxone should be administered in the event of a suspected opioid overdose.

## Warnings

- Medications kept at home must be locked in a safe place.
- Women should let their clinician know if they are pregnant or breast feeding.
- Combining methadone or buprenorphine with alcohol, sedatives, tranquilizers or other drugs that slow breathing can cause deadly side effects.

## Pain

- Pain can and should be effectively treated in patients with opioid addiction.
- For people who are not currently in treatment, methadone or buprenorphine can be effective for treating both opioid addiction and pain.
- The clinician may temporarily increase the patient's dose, or the frequency of dosing, of methadone or buprenorphine to effectively manage pain.
- The clinician may also prescribe additional pain-relieving medications or other treatments (such as physical therapy).
- Methadone or buprenorphine does not need to be stopped before surgery.
- If methadone or buprenorphine is stopped due to pain treatment, it should be restarted as quickly as possible.
- Naltrexone blocks the action of opioid pain medications. When needed, this blocking effect can often be overcome. In these instances, patients should be closely monitored in an emergency department or hospital setting.
- It is important that patients talk to their clinicians about their opioid addiction and work together to develop a pain management plan that both addresses the pain and minimizes their risk of a relapse.



## ASSESSMENT AND TREATMENT OF OPIOID USE DISORDER IN PREGNANT WOMEN

Untreated opioid use disorder can pose significant risk for pregnant women. It can lead to pregnancy complications as well as miscarriage, premature delivery, fetal growth restriction, and fetal death.

- Treatment for opioid use disorder should be initiated as early in the pregnancy as possible.
  - » Ideally the patient would be seen by a clinician experienced in pregnancy care as well as a clinician experienced in addiction treatment. However, if a specialist is not available, pregnant women should still seek prenatal care and discuss their need for opioid use disorder treatment.
- The clinician should complete a medical evaluation to identify any urgent medical issues that need to be addressed.
  - » Medical examination can include laboratory tests, physical examination, ultrasound, and questions about the patient's mental health and individual needs.
- The most important things that a pregnant woman with opioid use disorder can do are to get regular prenatal care and addiction treatment.
- Withdrawal can be dangerous during pregnancy. Patients should work with their clinician to manage withdrawal if they are going to stop or cut down on their opioid use (including prescription opioid use).
- The clinician should talk to the patient about the risks and benefits of the different treatment options. They should make a decision together based on what is best for the patient.
- Methadone and buprenorphine are the best treatments available for pregnant women with opioid use disorder. The clinician may recommend that the patient be hospitalized when the medication is started, especially in the third trimester. However, for most patients these medications can be safely started in an outpatient setting.



- The clinician may also recommend methadone or buprenorphine for pregnant women who are in recovery from opioid use disorder if the patient is at risk for relapse during pregnancy.
- Methadone or buprenorphine are the best options to treat opioid use disorder during pregnancy. However, if the patient is taking naltrexone prior to pregnancy and wants to continue it, the patient should discuss the potential risks with a clinician. Not enough is known about the safety of naltrexone during pregnancy.
- While methadone and buprenorphine have some similarities to other opioids like heroin or oxycodone, their specific properties make them much safer during pregnancy when used under the care of a clinician. These medications are longer acting, preventing opioid withdrawal, opioid overdoses, and the highs and lows that can harm the development of the fetus.
- Later in pregnancy the clinician may increase the dose of medication or have the patient take the medication more frequently to prevent cravings and support a more stable environment for the fetus.
- The clinician may recommend counseling or other behavioral treatments in addition to medication.
- Patients who are taking methadone or buprenorphine as prescribed, and are without other contraindications for breastfeeding, can and should breastfeed.
- After delivery the clinician will evaluate the patient to see if there is a need to adjust the dose of methadone or buprenorphine. Since there is a risk for relapse and overdose in the first year after delivery, patients should not stop taking these medications during this time.

# AN OPIOID OVERDOSE NEEDS IMMEDIATE MEDICAL ATTENTION

## Dos and Don'ts in Responding to Opioid Overdose

- **DO Call for Help (Dial 911)**
- **DO** support the person's breathing by administering oxygen or performing rescue breathing.
- **DO** administer naloxone (a drug that reverses the effect of opioids) as an injection or a nasal spray.
- **ALL FRIENDS/FAMILY SHOULD HAVE NALOXONE AND KNOW HOW TO USE IT.**
- **DO** put the person on their side if they are breathing independently.
- **DO** stay with the person and keep them warm.
- **DON'T** slap or try to forcefully stimulate the person—it will only cause further injury. If shouting, rubbing knuckles on the sternum (center of the chest or rib cage), or light pinching will not awaken the person, he or she may be unconscious.
- **DON'T** put the person into a cold bath or shower. This increases the risk of falling, drowning or going into shock.
- **DON'T** inject the person with any substance (salt water, milk, "speed," heroin, etc.). The only safe and appropriate treatment is naloxone.
- **DON'T** try to make the person vomit drugs that he or she may have swallowed. Choking or inhaling vomit into the lungs can cause death.

## QUESTIONS TO ASK YOUR CLINICIAN

- **What should I do if I think I have an opioid addiction?**
- **What types of treatment are available?**
- **What type of treatment would be best for me?**
- **How will my treatment change as I make progress in recovery?**
- **How will you make decisions about changes to my treatment plan, including changes to medications?**
- **What other services and supports are available to support my recovery?**
- **Are there differences in cost for treatment medications?**
- **Does insurance cover the cost of treatment medications?**
- **Where will I be treated?**



The American Society of Addiction Medicine represents leading addiction care providers dedicated to ensuring individuals suffering from addiction can access high quality care and enter into long term recovery.

For an online version of this patient guide and additional treatment and support resources, visit: [www.asam.org/guidelines](http://www.asam.org/guidelines)



<https://www.nj.gov/humanservices/dmhas/home/index.html>

Funding for this pocket guide made possible by the Substance Abuse and Mental Health Services Administration (SAMHSA) State Opioid Response Grants Catalogue of Federal Domestic Assistance (CFDA) No. 93.788

The views and opinions contained in this publication do not necessarily reflect those of SAMHSA or the U.S. Department of Health and Human Services, and should not be construed as such.



**ASAM** American Society of  
Addiction Medicine

ASAMPAT07203\_NJ